

Global Dimension in Categories of Diagrams

W. T. SPEARS

*Department of Mathematics, Kansas State University, Manhattan, Kansas 66502**Communicated by Saunders MacLane*

Received July 12, 1970

Let I be a finite partially ordered set considered as a category in the usual sense, and let \mathcal{O} be any abelian category. The relation of $\text{gl dim } \mathcal{O}^I$ (\mathcal{O}^I denotes the category of functors from I to \mathcal{O} and is usually referred to as the category of diagrams in \mathcal{O} over I) to $\text{gl dim } \mathcal{O}$ is the subject of a paper by Mitchell [3]. Among the questions he raises in his paper is the possible invariance of $\text{gl dim } \mathcal{O}^I - \text{gl dim } \mathcal{O}$ with respect to \mathcal{O} . This paper gives an example of a partially ordered set I with regard to which $\text{gl dim } \mathcal{O}^I - \text{gl dim } \mathcal{O}$ is dependent upon \mathcal{O} .

As Mitchell points out [2, pp. 229-233], the study of \mathcal{O}^I is one way to approach some problems in the global dimension of rings. Specifically, suppose E is a semigroup of matrix units, that is, $E \subset M_n$ where

$$M_n = \{e_{ij} \mid i, j = 1, \dots, n\} \cup \{0\},$$

and where the multiplication is that of matrix units: $e_{ij}e_{pq} = \delta_{jp}e_{iq}$. We may form the semigroup algebra over a field F , $F[E]$. Then we may ask what relation there is between the structure of E and the global dimension of $F[E]$. For example, if $E = M_n$, then the global dimension is zero; some other results along these lines have been obtained by Clark [1].

DEFINITION. \mathcal{O}^I will denote the category of all functors from the category I to the category \mathcal{O} . Alternatively, each functor may be thought of as a diagram in \mathcal{O} over I [2, p. 66]. For this reason, a category of functors \mathcal{O}^I is often referred to as a category of diagrams.

We will use the standard notations and conventions concerning \mathcal{O}^I that may be found in [2] or [3]. That is, for $D \in \mathcal{O}^I$ ($D: I \rightarrow \mathcal{O}$), $D(i)$ is denoted D_i ; for $i \leq j$, the morphism from D_i to D_j is denoted D_{ij} ; and if $f: D \rightarrow D'$ is a natural transformation, $f(i)$ is denoted f_i .

Let $T_i: \mathcal{O}^I \rightarrow \mathcal{O}$ be defined by $T_i(D) = D_i$ and $T_i(f) = f_i$ where $f: D \rightarrow D'$. A left adjoint for T_i , S_i , is defined by $S_i(A)_j = A$ for $i \leq j$ and is 0 otherwise; and $S_i(A)_{kl} = 0$ unless $i \leq k \leq l$ in which case it is 1_A .

The following result due to Mitchell [2, p. 228] is central to the study of the global dimension of \mathcal{O}' . Only the easily proved "if" part will be needed.

THEOREM 1. *An object D in \mathcal{O}' is projective if and only if $D \cong \bigoplus_{i \in I} S_i(P_i)$ where each P_i is a projective object in \mathcal{O} .*

The following result was obtained independently by Mitchell [3, p. 347] and the author.

DEFINITION. For each $i \in I$, define a functor $M_i : \mathcal{O} \rightarrow \mathcal{O}'$ by $M_i(A)_j = 0$ unless $j = i$ in which case it is A ; $M_i(A)_{kl} = 0$ for all $k, l \in I$.

The importance of the M_i 's as seen in the following result is that it allows the computation of $\text{gl dim } \mathcal{O}'$ from the homological dimension of members of \mathcal{O}' of a comparatively simple structure. This proof is due to the author.

THEOREM 2. *For any D in \mathcal{O}' , $\text{h.d. } D \leq \sup_{i \in I} \text{h.d. } M_i(D_i)$. Hence*

$$\text{gl dim } \mathcal{O}' = \sup\{\text{h.d. } M_i(A) \mid i \in I, A \in \mathcal{O}\}.$$

Proof. We will induct on the number of nonzero D_i 's in D . Certainly there is an i in I such that $D_i \neq 0$ and $D_j = 0$ if $j < i$. We have then the short exact sequence

$$0 \rightarrow D' \rightarrow D \rightarrow M_i(D_i) \rightarrow 0,$$

where $f : D \rightarrow M_i(D_i)$ is defined so that $f_i = 1_{D_i}$ and $f_j = 0$ for $j \neq i$, and where $D' \rightarrow D$ is the kernel of f .

From this exact sequence and an elementary result of homological algebra, we have that $\text{h.d. } D \leq \max(\text{h.d. } M_i(D_i), \text{h.d. } D')$. On the other hand, since $D_{j'} = D_j$ for $j \neq i$ and $D_{i'} = 0$, we have by induction that

$$\text{h.d. } D' \leq \max_{j \neq i} \text{h.d. } M_j(D_j).$$


Combination of these two results yields the desired inequality.

The following definition and result are from [3, pp. 345–346].

DEFINITION. The *rank* of an element of a partially ordered set is defined inductively. Minimal elements have rank 0. An element i has rank $\geq r$ if and only if it is preceded by two incompatible elements of rank $\geq r - 1$. The rank of i is r if $r \leq \text{rank } i \leq r + 1$.

PROPOSITION 3. *Let m be the maximum rank of an element of I and let $\text{h.d. } D_i \leq n$ for all i . Then $\text{h.d. } D \leq m + n$.*

Let I be the partially ordered set depicted in this Hasse diagram (directed upward).



THEOREM 4. Let \mathcal{F}_i denote the category of vector spaces over a field of characteristic i . Then $\text{gl dim } \mathcal{F}_2^I = 4$ and $\text{gl dim } \mathcal{F}_0^I = 3$.

Proof. It follows from Proposition 3 that $\text{h.d. } M_i(A) \leq 3$ for all $i \neq l$ and for all A in \mathcal{F}_2 or \mathcal{F}_0 . It will now be shown that for nonzero A , A in \mathcal{F}_2 implies that $\text{h.d. } M_l(A) = 4$, and A in \mathcal{F}_0 implies that $\text{h.d. } M_l(A) = 3$. This will be done by choosing a projective resolution for $M_l(A)$ without regard for the origin of A and then using Theorem 1.

Let $D^0 = S_l(A)$ which is projective by Theorem 1, and let $d^0: S_l(A) \rightarrow M_l(A)$ be the natural map. Now let K^0 denote $\ker d^0$ and define D^1 in terms of K^0 by letting $D^1 = \bigoplus_{i=1}^3 S_{a_i}(K_{a_i}^0)$ which is also projective. $d^1: D^1 \rightarrow K^0$ will be the epimorphism induced by the natural maps $S_{a_i}(K_{a_i}^0) \rightarrow K^0$. Then D^2 and $d^2: D^2 \rightarrow K^1 = \ker d^1$ are defined similarly. Let $D^2 = \bigoplus_{i=1}^6 S_{b_i}(K_{b_i}^1)$, and let d^2 be the epimorphism induced by the maps $S_{b_i}(K_{b_i}^1) \rightarrow K^1$. Let $K^2 = \ker d^2$.

Now it will be shown that if $A \in \mathcal{F}_2$, then K^2 is not projective; and if $A \in \mathcal{F}_0$, then K^2 is projective. First notice that $K_i^2 = 0$ for $i \neq c_1, \dots, c_4, m$, and that each $K_{c_i}^2$ may be regarded as a subobject of K_m^2 . Furthermore, a straightforward, if lengthy, calculation will verify that

$$K_{c_1}^2 + K_{c_2}^2 + K_{c_3}^2 = K_{c_1}^2 \oplus K_{c_2}^2 \oplus K_{c_3}^2.$$

(This is independent of A 's origin.)

Let B denote $(K_{c_1}^2 \oplus K_{c_2}^2 \oplus K_{c_3}^2) \cap K_{c_4}^2$. It follows by a straightforward calculation that if A is in \mathcal{F}_2 , then $B \neq 0$; and that if A is in \mathcal{F}_0 , then $B = 0$. Hence by Theorem 1, K^2 is not projective if A is in \mathcal{F}_2 and is projective if A is in \mathcal{F}_0 . Thus $\text{h.d. } M_l(A)$ equals 4 and 3, respectively, and it follows by Theorem 2 that $\text{gl dim } \mathcal{F}_2^I = 4$ and $\text{gl dim } \mathcal{F}_0^I = 3$.

ACKNOWLEDGMENT

The author expresses his gratitude for the help and guidance he received from his advisor, Edwin Clark. The author is also indebted to Barry Mitchell for his helpful and encouraging correspondence.

REFERENCES

1. W. EDWIN CLARK, Twisted matrix units semigroup algebras, *Duke Math. J.* **34** (1967), 417–424.
2. B. MITCHELL, "Theory of Categories," Academic Press, New York, 1965.
3. B. MITCHELL, On the dimension of objects and categories II. Finite ordered sets, *J. Algebra* **9** (1968), 341–368.